

7900056

TO ALL TO WHOM THESE PRESENTS SHAME COME;

## Seed Tec International, Inc.

Concreas, there has been presented to the

Secretary of Agriculture

AN APPLICATION REQUESTING A CERTIFICATE OF PROTECTION FOR AN ALLEGED NOVEL VARIETY of sexually reproduced plant, the NAME AND DESCRIPTION OF WHICH ARE CONTAINED IN THE APPLICATION AND EXHIBITS, A COPY OF WHICH IS HEREUNTO ANNEXED AND MADE A PART HEREOF, AND THE VARIOUS REQUIREMENTS OF LAW IN SUCH CASES MADE AND PROVIDED HAVE BEEN COMPLIED WITH, AND THE THERETO IS, FROM THE RECORDS OF THE PLANT VARIETY PROTECTION OFFICE; IN THE APPLICANT(S) INDICATED IN THE SAID COPY, AND WHEREAS, UPON DUE EXAMINATION MADE, THE SAID APPLICANT(S) IS (ARE) ADJUDGED TO BE ENTITLED TO A CERTIFICATE OF PLANT VARIETY PROTECTION UNDER THE LAW.

NOW, THEREFORE, THIS CERTIFICATE OF PLANT VARIETY PROTECTION IS TO GRANT UNTO THE SAID APPLICANT(S) AND THE SUCCESSORS, HEIRS OR ASSIGNS OF THE SAID APPLICANT(S) FOR THE TERM OF SEVENTEEN THAT FROM THE DATE OF THIS GRANT, SUBJECT TO THE PAYMENT OF THE REQUIRED FEES AND PERIODIC REPLENISHMENT OF VIABLE BASIC FEED OF THE VARIETY IN A PUBLIC REPOSITORY AS PROVIDED BY LAW, THE RIGHT TO EX-OTHERS FROM SELLING THE VARIETY, OR OFFERING IT FOR SALE, OR REPRODUCING IT, ORTING IT, OR EXPORTING IT, OR USING IT IN PRODUCING A HYBRID OR DIFFERENT THEREFROM, TO THE EXTENT PROVIDED BY THE PLANT VARIETY PROTECTION ACT \$42, AS AMENDED, 7 U.S.C. 2321 ET SEQ.)

SAFFLOWER

'S-742'

In Testimony Whereof, I have hereunto set my hand and caused the seal of the Mant Tariety Protection Office to be affixed at the City of Washington this 11th day of September

the year of our Lord one thousand nine hundred and eighty.


ΛDI	UNITED STATES DEPARTMEI AGRICULTURAL MARK LIVESTOCK, POULTRY, GRA PLICATION FOR PLANT VARIE	ETING SERVICE IN & SEED DIVISION		No certificate for pla		40-R3822 tion may
	RUCTIONS: See Reverse.	TY PROTECTION	N CERTIFICATE	be issued unless a co nas been received (5 t		tion form
1a.	TEMPORARY DESIGNATION OF VARIETY	1b. VARIETY NAMI		FOR OFFICI	AL USE ONLY	
	742-5-14 <sup>C</sup> -B-B	S-742		PV NUMBER 790003	56	
2.	KIND NAME	3. GENUS AND SPE	CIES NAME	FILING DATE	TH:00	A.M.
	Safflower	Carthamus t	inctorius L.	FEE RECEIVED	DATE	P.M.
4.	FAMILY NAME (BOTANICAL)	5. DATE OF DETER		s 500.00	3-5-7	19
	Compositae (asteraceae)	October 197	7.7	\$ 250.00	6/9/80	)
6.	NAME OF APPLICANT(S)	7. ADDRESS (Street	t and No. or R,F.D. No.,	City, State, and ZIP	8. TELEPHON	
	Pacific Oilseeds	Code) P.O. Box 10			(918)	
	Incorporated	Woodland, C	CA 95695		662-862	3
9.	IF THE NAMED APPLICANT IS NOT A PE ORGANIZATION: (Corporation, partnersh	HSON, FORM OF ip, association, etc.)	10. IF INCORPORAT DATE OF INCOR	ED, GIVE STATE AND PORATION	11. DATE OF PORATION	
	Corporation		California		1954	
12.	NAME AND MAILING ADDRESS OF APP ALL PAPERS:	LICANT REPRESENTA	ATIVE(S), IF ANY, TO	SERVE IN THIS APPLIC	ATION AND RE	CEIVE
	John W. Talbott, P.O. Box	1008, Woodland	l, CA 95695			
13.						
13.	I3A. Exhibit A, Origin and Bree		Voriety (See Section	52 of the Plant Variety	y Protection A	a <b>+</b> \
	13B. Exhibit B, Novelty Statem		variety (See Section .	02 Oj ine Fiant Variei	y Flotection A	-1.)
			· ·	<b>n</b> 4		
	X 13C. Exhibit C, Objective Descr			Plant Variety Protect	ion Office.)	
	☐ 13D. Exhibit D, Additional Des	cription of the Varie	ty.			
14a.	DOES THE APPLICANT(S) SPECIFY THAT SEED? (See Section 83(a). (If "Yes," answer			RIETY NAME ONLY AS	A CLASS OF C	ERTIFIED
14b.	DOES THE APPLICANT(S) SPECIFY THA LIMITED AS TO NUMBER OF GENERAT			IB, HOW MANY GENER BREEDER SEED?	ATIONS OF PRO	DDUC-
	YES NO	-	FOUNDATION	REGISTERED	CERTIFIE	D
15a.	DID THE APPLICANT(S) FILE FOR PROT name of countries and dates.)	ECTION OF THIS VAI	RIETY IN OTHER COU	NTRIES? YES	X NO (If ")	es," give
	name of countries and dates.)		en de la companya de Companya de la companya de la compa	rena de Maria de la composición de la c La composición de la	O Bylai	
			Asset San San San San			
15b,	HAVE RIGHTS BEEN GRANTED THIS VA	ARIETY IN OTHER CO	UNTRIES? YES	NO (If "Yes,"	give name of cou	intries
	and dates,)	· · · · · · · · · · · · · · · · · · ·				in Military (1977) January (1977)
	4.36					
	DOES THE ARRIVANTAN ACCES TO THE					· · ·
16.	DOES THE APPLICANT(S) AGREE TO TH JOURNAL? YES	NO A			<u>.</u>	
17.	The applicant(s) declare(s) that a viable replenished upon request in accordance	e sample of basic see e with such regulatio	d of this variety will l ns as may be applical	be furnished with the able.	application and	l will be
	The undersigned applicant(s) is (are) the variety is distinct, uniform, and stable 42 of the Plant Variety Act.	ne owner(s) of this se as required in Sectio	xually reproduced no n 41, and is entitled t	ovel plant variety, and to protection under th	believe(s) that e provisions of	the Section
4.	Applicant(s) is (are) informed that fals	e representation here	ein can jeopardize pro	otection and result in p	penalties.	
14	February 8, 1979		Johnso	Tellow	· · · · · · · · · · · · · · · · · · ·	<del></del>
. 144	(DATE)		John W. Ta	SIGNATURE OF APPLI 1bott, Executiv		esident

#### INSTRUCTIONS

GENERAL: Send an original copy of the application and exhibits, at least 2,500 viable seeds, and \$500 fee (\$250 filing fee and \$250 examination fee) to U.S. Dept. of Agriculture, Agricultural Marketing Service, Livestock, Poultry, Grain and Seed Division, Plant Variety Protection Office, National Agricultural Library Building, Beltsville, Maryland 20705. (See section 180.175 of the Regulations and Rules of Practice.) Retain one copy for your files. All items on the face of the form are self-explanatory unless noted below.

#### ITEM

- Give the date the applicant determined that he had a new variety based on (1) the definition in section 41(a) of the Act and (2) the date a decision was made to increase the seed.
- Give: (1) the genealogy, including public and commercial varieties, lines, or clones used, and the breeding method; (2) the details of subsequent stages of selection and multiplication; (3) the type and frequency of variants during reproduction and multiplication and state how these variants may be identified and (4) evidence of uniformity and stability.



- Give a summary statement of the variety's novelty. Clearly state how this novel variety may be distinguished from all other varieties in the same crop. If the new variety most closely resembles one or a group of related varieties:

  (1) identify these varieties and state all differences objectively; (2) attach statistical data for characters expressed numerically and demonstrate that these differences are significant; and (3) submit, if helpful, seed and plant specimens or photographs of seed and plant comparisons clearly indicating novelty.
- Fill in the Exhibit C, Objective Description form, for all characteristics for which you have adequate data.
- Describe any additional characteristics that are not described, or whose description cannot be accurately conveyed in Exhibit C. Use comparative varieties as is necessary to reveal more accurately the description of characteristics that are difficult to describe, such as, plant habit, plant color, disease resistance, etc.
- If "YES" is specified (seed of this variety be sold by variety name only as a class of certified seed) the applicant may NOT reverse his affirmative decision after the variety has either been sold and so labeled, his decision published, or the certificate has been issued. However, if the applicant specified "NO," he may change his choice. (See section 180.16 of the Regulations and Rules of Practice.)
- See section 42 of the Plant Variety Protection Act and section 180.7 of the Regulations and Rules of Practice.

EXHIBIT A: ORIGIN AND BREEDING HISTORY S-742.

S-742 was developed by pedigree selection from the following multiple cross completed in 1970:

(Ul421 x Gila) x POI-261 bc<sup>1</sup>) x UC1 x (POI-261 x F64-1-47) x PCOY

U1421--a rust resistant line acquired in 1960 from Utah State University, Logan, Utah.

POI-261--a high oil striped pericarp breeding line developed by Pacific Oilseeds Incorporated.

F64-l-47-a fusarium resistant selection acquired in 1965 from C. A. Thomas USDA Plant Pathologist. This selection by Dr. Thomas was from the cross: (N6 x Gila bc $^{\rm l}$ ) x POI-414. POI-414 is a high yielding breeding line developed by Pacific Oilseeds Incorporated.

PCOY--a rust resistant breeding line released by Mr. A. L. Urie, USDA Agronomist, Utah State University.

In 1972,  $F_2$  segregants of the above cross were evaluated for general appearance and rust resistance. Rust resistant  $F_3$  progeny were grown in the Yolo bypass near Woodland, California. Row number 742 was rated as 90% resistant to fusarium wilt. Selfed seed of rust and fusarium resistant selection 742-5 was grown in 1974 again in the Yolo bypass. Rust was rated as resistant and fusarium incidence was only three percent. The entire row was caged and selfed seed from plant 742-5-14°C having 46.5% oil measured by standardized NMR analysis was advanced to the  $F_6$  generation. Selfed seed from 742-5-14°C was evaluated for resistance to verticillium wilt during 1975-1976 by Dr. C. A. Thomas, USDA Plant Pathologist. Selection 742-5-14°C was rated 100% resistant to the cotton defoliant isolant of verticillium. This compared to 100% resistance in the USDA line VFR and 100% susceptibility in the variety 'Gila'.

In 1976 742-5-14° was grown in rows and appeared uniformly homozygous. The line was bulked and entered in POI yield trials in 1977. Results from 1977 yield trials warranted increase of breeders seed in Mexico in 1977-1978. Seed returned from increase in Mexico was planted in 1978 in the Sacramento Valley and increased in isolation for commercial seed production in 1979. This line has been tentatively named S-742.

Results from 1977 yield trials warranted in 1977-1978. Seed returned from increase in Mexico was planted in 1978 in the Sacramento Valley and increased in isolation for commercial seed production in 1979. This line has been tentatively named S-742.

Results from 1977 yield trials in 1977-1978. Seed returned from increase in Mexico was planted in 1978 in the Sacramento Valley and increased in isolation for commercial seed production in 1979. This line has been tentatively named S-742.

Results from 1977-1978. Seed returned from increase in Mexico was planted in 1978 in the Sacramento Valley and increased in isolation for commercial seed production in 1979. This line has been tentatively named S-742.

TH:tb 2/8/79

# ACIFIC OILSEEDS INCORPORATED

Safoa SAFFLOWER • HYBRID SORGHUM • HYBRID SUNFLOWER

916 - 662-8623

August 24, 1979

P. O. BOX 1008 WOODLAND, CA 95695

Mr. Joseph T. Higgins
Examiner, Plant Variety
Protection Office
USDA
National Agricultural Library Bldg.
Bettsville, MD 20705

Dear Mr. Higgins:

I have enclosed supportive data and descriptions on safflower varieties S-541 and S-742 (application numbers 7900056 and 7900057). The differences between presently protected varieties (S-317 and S-400), standard varieties, S-541 and S-742, are summarized in table form and evidenced by accompanying photographs.

In your letter of April, 1979, you asked about the reddish-orange dry down color of S-742. This is a darker orange than that of "Gila" and is unique. To my knowledge, no other varieties on the market have this shade of dry down floret. I believe the color shows up somewhat on the photographs.

With respect to determining the stability of the S-541 and S-742 genotypes, this was done by bulking advanced generation plants that were identical in flower colors, height, plant morphology, pericarp type, and oil type. No off-type plants were noted in either seed increase nor have off-type plants been noted in yield trials conducted throughout California.

Also, to the best of my knowledge, the variety "Carmex" is tested and sold only under that designation.

I believe the data presented are as accurate as possible. Of course, heights and oil contents change somewhat according to planting dates and available moisture, but, in general, differences remain relative and are apparent.

I hope you find our application is in order. Many thanks for your patience as we were awaiting the collection of materials.

Sincerely,

PACIFIC OILSEEDS INCORPORATED

Thomas C. Heaton Research Director

TCH/cbs

ತ

#### PACIFIC OILSEEDS INCORPORATED

EXHIBIT B: NOVELTY STATEMENT ON S-742.

Safflower variety S-742 is a linoleic acid oil type cultivar. Variety S-742 is Pacific Oilseeds Incorporated's first triple resistant safflower variety. S-742 has shown good field resistance to the prevalent race 2 of Fusarium oxysporum and has performed well in areas where the new race 4 is suspected to occur. Resistance to the severe strain of verticillium has been demonstrated in USDA seedling screening tests. Incorporation of the rust resistance gene PCOY (a gene originally transferred from the wild species Carthamus oxyacantha) makes S-742 resistant to all strains of safflower rust, Puccinia cartami. However, it appears that a new strain of rust in localized areas may be developing. S-742 is not immune to this strain, although the strain has not become important. If this is a new strain, resistance has not yet been identified in either public or private materials.

4/28/80

S-742 is distinguished from other varieties by its large head diameter, its abundance of lanceolate bracts, its reddish-orange dry down color, and its triple disease resistance. S-742 most closely resembles the related varieties "Gila" and "Frio" in plant habit and branching pattern, but is easily distinguished by its darker reddish-orange dry down, higher oil content (striped pericarp), larger head diameter, and numerous head bracts. S-742 is completely unique from standard safflower varieties. Table 6 compares and contrasts S-742 with seven other common safflower varieties.

Yield results for 1976 through 1978 are presented in Tables 1 through 3. S-742 was not entered in USDA trials due to a limit on the number of entries that could be made. It can be seen that S-742 is an improvement over the two main high linoleic acid oil type commercial varieties in California, S-208 and S-400. S-742 combines high oil content with competitive yields and excellent disease resistance. We expect S-742 to be popular in areas where rust and wilt diseases are detrimental to safflower production.

TCH/cbs 8/23/79

TABLE 6: COMPARISON OF SAFFLOWER VARIETAL CHARACTERISTICS

· I									
FIRST FLOWER (DAYS)	16	06	82	87	87	96	35	16	
VERTICILLIUM	<b>-</b>	œ	νı	ν	S	⊢	·	۰	
FUSARIUM WILT (RACE 2)	oc.	œ	S	S	Ś	S	œ	≃ .	
RUST d	S	œ	· vs	s	s,	S	S	<b>v</b>	
OIL CONTENT C (%)	46.7	47.1	40.1	41.2	42.5	43.6	43.6	44.1	
PERICARP	gray	purple striped	white	white	white	white	gray striped	gray striped	
O1L TYPE	linoleic	linoleic	linoleic	linoleic	linoleic	linoleic	linoleic	oleic	
FLOWER COLOR (DRY)	orange	reddish orange	orange	yellow	orange	orange	yellow	yellow	
FLOWER COLOR (BLOOM)	yellow	yelłow	yellow	yellow	yellow	yellow	yellow	yellow	
NUMBER MAIN BRANCHES	9 ا		5 -	٠. د د	9	7	9-	ا ئ	
HEAD H DIAMETER (CM) B	2.6 0.2	3.2	2.6 0.2	2.4	2.7	2.5	2.6 0.2	2.6	
TH ½ LENGTH IS BRACT WIDTH DI (CM)	0.9 0.1	0.1	1.2	 	0.1	0.0 0.1	1.0	0.8	
LENG BRAC (CM)	2.8	3.5	3.0	3.4	3.0	3.0	3.1 0.3	2.9	
NUMBER <u>5</u> BRACTS	20	33	18 2	17	20	9 -	25 3	24	
PLANT A HEIGHT (CM)	8 8	79	74 2	79	83 <b>2</b>	ا 2	8 2	81	
1	i×ω	ıx <b>ı</b> o	ı×ω	ı×ιο	ı×ıo	i×ĸ	αν×ι	ıxıo	
VARIETY	5-541	s-742	Gila	US-10	Frio	S-208	S-400	5-317	

 $\bar{x}$  = average of 20 observations;  $\delta$  = standard deviation.

all bract and head measurements were taken on primary heads.

oil contents from the same location/same year.

T = tolerant; R = resistant; S = susceptible. ाज ।

TH/cs

EXHIBIT B: Yield Data on S-742 (Percent of oil by NMR)

Table 1. Advanced Safflower 1976-2 River Garden Farms, California

Entry	Yield lb/A	Oil %	Test wt lb/bu	Oil/A	Days to 100% bloom	Height cm
742-5-14 <sup>C</sup> -B	3621	48.3	41.1	1749	80	68
S-400	3588	44.4	42.2	1593	79	68
S-296	3416	43.3	42.4	1479	79	71
S-317	3204	. 42.0	42.9	1346	78	81
S-208	3058	43.1	43.0	1318	76	76
Erhol	2788	46.7	43.9	1302	77	71
				-		
Table 2. Adva	anced Safflow	ver 1977-8	Clarksburg, (	California		
742-5-14 <sup>С</sup> -В <b>-</b> В	4018	47.2	41.9	1896	98	114
S-208	3660	44.4	43.2	1625	96	113
s-400	3463	45.2	41.9	1565	98	122
s-296	3399	44.0	42.4	1495	98	122
Table 3. Adva	unced Safflow	er 1978-8 (	Clarksburg, Ca	alifornia		
s-317	4178	43.4	41.5	1813	95	114
US-10	3453	39.1	43.0	1350	89	107
S-541	3342	46.2	41.9	1544	92	114
s-208	3233	42.8	41.7	1384	89	107
742-5-14 <sup>C</sup> -B <b>-</b> B	3209	46.0	40.1	1476	91	107
S-400	3111	43.6	41.0	1357	92	107
s-296	3083	42.2	40.8	1301	95	114
Gila	3069	41.1	41.3	1261	89	91
Frio	2938	42.7	40.0	1255	89	107
	•					

TH:tb 2/8/79



#### PACIFIC OILSEEDS INCORPORATED

Table 4.

### 7900056

Variety	Hull	Protein %	<b>୍ତୀ</b> ୫୫	WIJS Todine Value
S-208	34.6	12.7	41.7	145.8
S-400	32.1	13.0	43.4	143.8
s-541	30.1	14.9	44.2	141.9
s-742	31.8	13.1	45.7	141.8

UNDECORTICATED SEED ANALYSIS\*

## FATTY ACID DISTRIBUTION ANALYSIS\*

Variety	Carbon No.	14	16	16:1	18	18:1	18:2	20	20:1	22
s-208	8	.1	6.6	.1	2.3	11.1	79.2	.2	.2	.2
S-400	2.3	.1	6.8	.1	3.0	12.0	77.5	.2	.2	.2
S-541		.1	7.2	.1	2.8	14.0	75.0	.4	.3	.2
S-742		.1	7.4	.1	3.2	12.6	75.8	.4	.2	. 2

	Acids Saturated	Acids Unsaturated	Oleic Acid	Linoleic* Acid
Variety	*	.8	*	<b>%</b>
<b>S-20</b> 8	9.4	90.6	11.1	79.2
S-400	10.3	89.7	12.0	77.5
S-541	10.7	89.3	14.0	75.0
S-742	11.3	88.7	12.6	75.8

<sup>\*</sup>Samples from 1978 Yield trial at River Garden Farms. Analyses done by PVO/Richmond Mr. John Wood, December 1978.

1/4/79 tb FORM GR-470-22 (2-15-73)

## AGRICULTURAL MARKETING SERVICE GRAIN DIVISION HYATTSVILLE, MARYLAND 20782

EXHIBIT C (Safflower)

### **OBJECTIVE DESCRIPTION OF VARIETY**

INSTRUCTIONS: See Reverse. SAF	FLOWER (CARTHAMUS TINCT		
PACIFIC OILSEEDS INCORPORATED		FOR OFFICE	AL USE ONLY
ADDRESS (Street and No. or R.F.D. No., City, State,	and ZIP Code)	79000	56
P.O. Box 1008		VARIETY NAME OR T	
Woodland, CA 95695		3-74	2
Place the appropriate number that describes the Place a zero in first box (e.g. 0 8 9 or 0			
1. MATURITY (From Emergence):			
Location: 1 = CALIFORNIA & ARIZONA	2 = MIDWEST	1 2 0 NO. C	OF DAYS TO MATURIT
0 0 NO. OF DAYS EARLIER THAN 0 0 2 NO. OF DAYS LATER THAN 3	1 = GILA 2 = FRIO	3 = US-10	
2 Maturity Class: 1 = EARLY (less than 1 3 = MEDIUM LATE (12	· · · · · · · · · · · · · · · · · · ·		
2. PLANT HEIGHT AT MATURITY:		—— \	
1 0 7 CM. HEIGHT 0	0 CM. SHORTER THAN	2	2 = FRIO 3 = US-1
3. FLOWER COLOR:    Type: 01 =   Fresh Flower   WHITE   WHITE	GREYISH-WHITE 05 ORANGE BASE 05 YELLOW (US-10) LIGHT ORANGE BASE 10	Fresh Flower  = ORANGE  = REDDISH ORANGE  = YELLOW BASE & TIPS OF LOBES ORANGE  = PALE-YELLOW  = OTHER (Specify) yellow	·· DEEP RED · ORANGE · PALE-YELLOW
4. SPINES ON INVOLUCRAL BRACTS:			
2 1 = ABSENT 2 = PRESENT			e de la companya de l
3 Location: 1 = TIP ONLY 2 = TIP & FI	EW BASAL 3 = TIP & ALONG M	ARGINS 4 = MARGINS O	NLY
0 3. MM. LENGTH (A)	1 2 NUMBER (B)	0 3 6 SPIN	E INDEX (A. x B)
2 Spine Index Class: 1 = 0 -20 2 = 21-40	3 = 41-60 4 = 61-80 5 =	81-100 6= 101-120	
5. HEADS (For Plant Populations of 593,000 Plants	/Hectare):		
0 2 5 MM. DIAMETER (Primary Heads)	TAGE A		
Seed Shattering Percentage: 1 = 1-10	2 = 11-30 3 = OVER 30		
6. SEED:			
Colors	BEAM 3 = GRAY URPLE STRIPED 7 = BROWN STRI	4 = GRAY WITH WI PED 8 = BROWN BLOTC	
3 Hull Type: 1 = NORMAL 2 = THIN-H	ULCED 3 = STRIPED 4 = RE	DUCED	
0 3 MM. WIDTH	0 6 MM. LENGTH	0 2 6 GRA	MS PER 1000 SEED
7. SEEDLING VIGOR: (6 weeks after seeding at 2.	5 cm, depth with ample moisture for g	ermination; mean of 20 plants)	- /
1 2 NUMBER OF NODES	0 1 2 CM. TALL (Soil Surface	to Tip)	

	·		•		790	0056	5-742
FORM GR-470-22 (REV	·····				<u></u>		
8. COLD RESISTANCE	AT DIFFERENT STA	GES AND TEMPER	ATURES:				
3 Rosette:						-	,
4 Bolting:	1 = -10°C. 2 =	-5°C. 3 ≈ 0°C.	4 = 5° C.	5 = 10° C.			
5 Flowering:							
9. DISEASE: (0 = No	t Tested; 1 = Susceptil	le: 2 = Resistant)				·	
2 RUST (Specify rac resistant	•		THORA ROO	т вот	PYTHIUM	ROOT ROT	
2 FUSARIUM WILT		2 VERTICIL	LIUM WILT		O CERCOSP	ORA LEAF SI	тот
0 SCLEROTINA ST	EM ROT	O ALTERNA	RIA LEAF S	РОТ	O ALTERNA	RIA BUD RO	Т
0 BOTRYTIS HEAD	ROT	0 RHIZOCTO	NIA BLIGH	г	0 BACTERIA	AL BLIGHT	•
O CUCUMBER MOS	AIC	O PHYLLOD	<b>7</b> .		O OTHER (S	pecify)	
10. INSECT AND NEM	ATODE: (0 = Not T	ested; 1 = Susceptib	le; 2 = Resist	ant)			
O GREEN PEACH A	PHID	0 LEAF-CUR	L PLUM APH	HID	0 BLACK BE	AN APHID	•
0 WESTERN FLOWE	R THRIPS	0 LYGUS BU	GS		0 STINKBU	3S	
O ROOT-KNOT NEM	IATODE	O OTHER (Sp	ecify)		 -		
11. INDICATE A VARI	ETY THAT MOST CLO	SELY RESEMBLES	THAT SUBI	WITTED:			
CHARACTER	VA	RIETY	С	HARACTER		VARIETY	
Frost Hardiness	s-400		L	odging	S-317		
Seed Shattering	FR10		N	o, of Branches	FR10		
Seedling Vigor	FR10						
12. GIVE THE FOLLO	WING DATA FOR SU	BMITTED AND A SI	MILAR VAR	IETY *:			
VARIETY	HULL (%)	PROTEIN (%)	OIL (%)	IODINE (%)	ACIDS SATURATED (%)	ACIDS UNS	LINOLEIC (%)
Submitted	21 0	72.1	<i>1</i>	141 0	- 77 2	12.6	75.8

		PROTEIN (%)	OIL (%)	IODINE	ACIDS	ACIDS UNSATURATED		
VARIETY	HULL (%)			(%)	SATURATED (%)	OLEIC (%)	LINOLEIC (%)	
Submitted	31.8	13.1	45.7	141.8	11.3	12.6	75.8	
Similar	32.1	13.0	43.4	143.8	10.3	12.0	77.5	
Name of Similar Variety	S-400	> 80l	7 4/15/80					

<sup>\*</sup>Hull, protein, and oil percentages expressed for whole undecorticated seed; acids expressed as percentages of oil.

#### REFERENCES

- 1. Knowles, P.F. & M.D. Miller. 1965. Safflower. Cal. Ag. Exp. Sta. Circ. 532. 51 p.
- 2. Weiss, E.A. 1971. Castor, Sesame, and Safflower. Barnes & Noble, Inc. N.Y. 901 p.

Nickerson's or any recognized color fan may be used to determine plant colors of described variety.

COMMENTS: